# privet Prebiotic 500 g | 500 g




| | |
|---|---|
| **Manufacturer** | priVet Farming® |
| **Product number** | 4855 |
| **Weight** | 0.35kg |
| **Length** | 98mm |
| **Width** | 98mm |
| **Height** | 124mm |

## Product description

***Complementary feed (powder) for ruminants, calves, pigs, and poultry***

Nutritional-physiological support of the intestinal flora

In a healthy intestine the balance of the intestinal flora is intact. Pathogenic bacteria thus do not find an ideal basis for life and can only develop with great difficulty. Since a large part of the immune system is located in the intestine, an intact intestinal flora favors a well-functioning immune system.

**privet Prebiotic** is a feed complement with a special mineral complex that has toxin-binding properties. It was specifically designed for particularly intense, nutritional-physiological support in digestive disorders. Due to the feed composition, fattening poultry may suffer from digestive disorders, which are indicated by thin feces, for example. **privet Prebiotic** supports the intestine especially in the binding power of liquids - important for animals prone to indigestion. The development of a healthy intestinal flora is essential for a healthy development of young animals.

**Composition:** linseed expeller, black cumin seed expeller

**Additives/kg:** Technological additives: bentonite (1m558i) 100 g.
The total amount of bentonite must not exceed the permitted maximum level in complete feedingstuff of 20000 mg/kg of complete feedingstuff.

**Analytical constituents:** crude protein 15.5%, crude fat 4.6%, crude fiber 5.2%, crude ash 16%, ash insoluble in HCl 14%, calcium 0.61%, phosphor 0.23%, magnesium 0.38%, sodium 0.27%, lysine 0.59%, methionine 0.26%

**Feeding recommendation:** ***Cattle/pigs:*** 0,5-2 g/kg feed. ***Poultry:*** 1 g/kg feed. ***Calves:*** stir 5-20 g/aminal into the milk. 1 tsp. corresponds to approx. 3.5 g / 1 tbsp. corresponds to approx. 11 g.

The simultaneous oral use with macrolides shall be avoided. The simultaneous use with robenidine shall be avoided. The simultaneous use with coccidiostats other than robenidine is contraindicated with level of bentonite above 5000 mg/kg of complete feed.

## Additional information

**PZN DE** 19680440